

Eliminating Sex Disparities in Dyslipidaemia Screening, Diagnosis and Treatment

Amelia E Power¹ and Dinesh K Kalra²

1. Department of Medicine, University of Louisville School of Medicine, Louisville, KY, USA; 2. Cardiovascular Medicine, Rudd Heart & Lung Center, University of Louisville School of Medicine, Louisville, KY, USA

<https://doi.org/10.17925/HI.2025.19.2.2>

Despite strides in cardiovascular disease (CVD) management, dyslipidaemia remains a significant yet underdiagnosed and undertreated risk factor, particularly among women. Sex-based disparities persist in screening, diagnosis and treatment, leading to suboptimal management and increased CVD risk in female populations. This article explores the current literature on sex disparities in dyslipidaemia, analysing screening guidelines, diagnosis trends and treatment gaps. It examines factors influencing lipid metabolism across a woman's lifespan, including hormonal fluctuations, pregnancy, menopause and their impact on CVD risk. The article also highlights barriers to effective lipid management in women, including clinician biases, inadequate screening and lower prescription rates of statin and non-statin therapies. Women are less likely to undergo lipid screening despite having significant CVD risk factors. Even when diagnosed, they receive statin therapy at lower rates than men, and treatment intensification is less frequent. Additionally, clinical trials assessing lipid-lowering therapies often underrepresent women, limiting the applicability of evidence-based recommendations. The lack of sex-specific risk assessment tools further contributes to missed opportunities for prevention and treatment. Addressing disparities in dyslipidaemia management is crucial to reducing the burden of CVD in women. Enhancing awareness among clinicians, improving screening strategies, incorporating sex-specific risk factors into predictive models and increasing female representation in clinical trials are essential steps towards equitable cardiovascular care.

Keywords

Cardiovascular disease, diagnosis, dyslipidaemia, hyperlipidaemia, lipid metabolism, screening, sex disparities, statin, treatment

Disclosures: Amelia E Power and Dinesh K Kalra have no financial or non-financial relationships or activities to declare in relation to this article.

Review process: Double-blind peer review.

Compliance with ethics: This article involves a review of the literature and did not involve any studies with human or animal subjects performed by any of the authors.

Data availability: Data sharing is not applicable to this article as no datasets were generated or analysed during the writing of this article.

Authorship: All named authors meet the criteria of the International Committee of Medical Journal Editors for authorship for this manuscript, take responsibility for the integrity of the work as a whole and have given final approval for the version to be published.

Access: This article is freely accessible at touchCARDIO.com. © Touch Medical Media 2025.

Received: 7 May 2025

Accepted: 5 August 2025

Published online: 6 October 2025

Citation: *Heart International*. 2025;19(2):Online ahead of journal publication

Corresponding author: Dinesh K Kalra, 201 Abraham Flexner Way, Suite 600, Louisville, KY 40202, USA. E: dinesh.kalra@louisville.edu

Support: No funding was received in the publication of this article.

Despite advancements in mortality rates from cardiovascular disease (CVD) over the last 50 years, it remains the leading cause of death for women and the third leading cause for women aged 18 to 44.¹ A 2011 report from the American Heart Association (AHA) starkly states that one in every two women will die from a cardiovascular-related disorder.² With the increasing prevalence of metabolic syndrome and obesity, the urgency of focusing on the early identification of dyslipidaemia to enhance the primary prevention of CVD cannot be overstated.

Women have been found to have an increasing prevalence and incidence of dyslipidaemia, with an estimated 40.4% of women having a total cholesterol (TC) level >200 mg/dL.³ Dyslipidaemia and the development of atherosclerotic cardiovascular disease (ASCVD) in women are associated with the highest population-attributable risk for myocardial infarction (MI) when compared to other CVD risk factors such as smoking, diabetes or obesity.⁴ However, although >90% of women have at least one CVD risk factor defined by the AHA and the United States Preventive Services Task Force (USPSTF), as many as 50% of this population are not screened appropriately with surveillance lipid panels.⁵ Even when women are screened, the female sex is associated with a lower likelihood of receiving statin therapy compared to the male sex, suggesting that many women with dyslipidaemia are not being diagnosed and treated appropriately.⁶ Women are less likely to be prescribed high-intensity statin therapy and achieve low-density lipoprotein cholesterol (LDL-C) goals, despite having risk factors. They are also less likely to receive non-statin lipid-modifying medications.^{7,8} The potential consequences of these disparities are severe, including increased mortality rates and reduced quality of life. In this article, we highlight disparities in the screening, diagnosis and treatment of women with dyslipidaemia. We also provide a call to action to continue evidence-based treatment of dyslipidaemia and focus on strategies to advance women's cardiovascular health.

A brief overview of the physiology of lifetime lipids in women

Female physiology causes more significant fluctuations of lipid levels throughout the lifespan. Before adolescence and early adulthood, lipid levels are generally similar between men and women, except for fluctuations associated with the menstrual cycle and pregnancy, which can cause an increase in lipid levels of up to 50%.⁹ Men generally have a shorter time to peak cholesterol development (45–49 years), while women reach peak cholesterol about 10 years later in their mid-50s.¹⁰

In adolescence, paediatricians perform universal screening lipid panels on children who are thought to be at risk for metabolic syndrome. These panels are usually obtained when the children are 9–11 years old and 17–21 years old to identify higher-risk individuals, including those with genetic disorders such as familial hyperlipidaemia.^{11,12} If a child is found to have high lipid levels, the first recommendation is lifestyle modification, followed by statin therapy. As mentioned above, fluctuations in lipid levels are associated with the menstrual cycle; therefore, clinicians must consider this when screening adolescent females for lipid disorders. Normal lipid panel variation is estimated at 5–8%. TC and LDL-C levels rapidly increase after menses and peak in the follicular phase before declining throughout the luteal phase. However, meta-analysis of triglyceride (TG) levels during the menstrual cycle has found that levels do not fluctuate a clinically significant amount. The rise and peak of TC and LDL-C correspond to the oestrogen peak during the menstrual cycle, whereas the decline corresponds to the progesterone peak.¹³

Many teenage girls will initiate contraception during the years they are being screened for dyslipidaemia. The National Lipid Association (NLA), a leading authority in lipid management, has found that specific birth control methods, such as oral contraceptive pills (OCPs), can cause an increase in TG levels, particularly those formulations of OCPs that have higher concentrations of oestrogen.¹⁴ There are currently no guidelines on when to screen women during the menstrual cycle, but a suggestion is to compare like samples. Of note, as metabolic syndrome increases in this young population, a higher proportion of women have been diagnosed with polycystic ovarian syndrome (PCOS), affecting an estimated 5–13% of the population.¹⁵ This syndrome is characterized by menstrual irregularity or amenorrhoea, hyperandrogenism, insulin resistance and polycystic ovaries. PCOS can be associated with dyslipidaemia and increased CVD risk; these patients should be counselled on lifestyle management and have aggressive screening for dyslipidaemia, with international guidelines recommending every 6–12 months.^{16–18}

Transitioning from adolescence into adulthood, new factors influence lipid levels in women, particularly pregnancy. Several studies have found that unfavourable CVD risk factors in the pre-pregnancy period, including dyslipidaemia, are associated with an increased risk of developing pregnancy complications such as pre-eclampsia.¹⁹ An analysis of patients in the Framingham Heart Study analysed over 500 parent–offspring pairs after measuring LDL-C before the offspring's birth. It concluded that high maternal pre-pregnancy LDL-C levels are associated with high adult offspring LDL-C levels, even when accounting for lifestyle, anthropometric and inherited factors, suggesting possible epigenetic changes.²⁰ In an analysis of data from the National Health and Nutrition Examination Surveys from 1999 to 2014 of women aged 20–44, only 39% of pregnant women had ideal cholesterol levels, suggesting that pre-pregnancy diagnosis and treatment of dyslipidaemia may miss many women who would benefit from it.²¹

Women experience increases in TC and TGs during pregnancy, peaking at term delivery. TC is estimated to increase between 35 and 37% in uncomplicated pregnancies. TGs increase between two- and three-times baseline levels.²² Higher levels of TC, TG, LDL-C and lower high-density lipoprotein cholesterol (HDL-C) have been associated with adverse outcomes, such as preterm birth, gestational diabetes and pulmonary embolism.²³ There are no screening guidelines for dyslipidaemia in pregnant women. Screening before pregnancy is considered the most beneficial, which is the current recommendation of the NLA.²⁴ If abnormal values are found, they should be monitored during pregnancy and followed up at 6 weeks postpartum.²¹ In a *Journal of the American*

College of Cardiology (JACC) update on recommendations for primary prevention in women, it is recommended that women experiencing adverse pregnancy outcomes (APOs) such as gestational hypertension (HTN), gestational diabetes, intrauterine growth restriction, placental abruption, obesity, sleep disorders and maternal age >40 years undergo cardiovascular screening within 3 months of postpartum.¹⁸


Until recently, there were limited effective options to treat dyslipidaemia during pregnancy due to the contraindication label of statins by the Food and Drug Administration (FDA).²⁵ In 2021, the FDA requested the removal of that classification, and the current recommendation is that treatment decisions should be made individually, based on the patient's risk profile, and through shared decision-making between the patient and clinician.²⁶ According to the NLA, clinicians can consider other therapies, such as bile acid sequestrant colesevelam and LDL-C apheresis in extreme cases.²⁷ Ezetimibe has been associated with adverse foetal effects in animal studies and is not recommended, and the safety or efficacy of PCSK9 inhibitors has not been determined in pregnancy.²⁷ In rare cases, such as gestational hypertriglyceridaemia-induced pancreatitis, it might be necessary to perform therapeutic plasma exchange, as there is significant morbidity and mortality associated with this condition.²⁸

After menopause, when women lose the mediating effects of oestrogen, their overall lipid levels tend to be higher compared to men, increasing their risk of CVD during the latter half of life.²⁹ Menopause is associated with an atherogenic lipid profile with increased LDL-C and TC and a decline in HDL-C levels. In one meta-analysis, women experienced an annual increase of 1.5 mg/dL and 1.2 mg/dL in TC and LDL-C levels, respectively, following menopause.³⁰ Certain factors have been linked to early menopause, such as smoking, autoimmune disease, low socioeconomic status and Black or Hispanic race, suggesting that these groups should have more aggressive screening for dyslipidaemia, given that they are at increased risk for cardiovascular morbidity and mortality.^{31,32} Menopause has also been associated with an increase in waist circumference, increased central adiposity, loss of skeletal muscle and an increase in fat mass due to oestrogen deficiency, which in turn increases CVD risk.³³ The factors influencing cholesterol in women across their lifespans are illustrated in *Figure 1*.^{21,34}

Factors influencing dyslipidaemia change throughout a woman's life, illustrating how crucial proactive screening is for reducing future CVD risk. However, as discussed in the next section, younger women rarely undergo screening, even with increasing risk factors in this population, such as obesity, insulin resistance and metabolic syndrome. It is the authors opinion that most women are currently being under-screened for dyslipidaemia.

Sex disparities in screening

The AHA and USPSTF publish recommendations for screening for dyslipidaemia, which are outlined in the following paragraph. However, many of these guidelines recommend different ages for initial screening based on the presence or absence of CVD risk factors, creating confusion. The AHA recommends screening adults older than 20 who are not on lipid-lowering therapy to document their baseline ASCVD risk and LDL-C levels.³⁴ The USPSTF updated its guidelines in 2022 to recommend screening all adults aged 40–75 with one or more cardiovascular risk factors and treating them with a statin if a 10-year CVD risk is 10% or greater based on pooled cohort equations (PCEs).³⁵ Before these updated guidelines, the USPSTF recommended screening men at 35 years old and women at 45 for dyslipidaemia.³⁶

Figure 1: Physiology of lipids across the female lifespan and factors influencing lipid levels^{21,34}

FH = familial hyperlipidaemia; APOs = adverse pregnancy outcomes; ASCVD = atherosclerotic cardiovascular disease; DM = diabetes; HDL-C = high-density lipoprotein cholesterol; LDL-C = low-density lipoprotein cholesterol; TC = total cholesterol; TG = triglyceride.

The presence of CVD risk factors should prompt earlier screening by health professionals for women and men alike. These factors have been modelled in PCEs, such as the 10-year ASCVD risk calculator, which takes into account a patient's age, sex, race, systolic and diastolic blood pressure, cholesterol panel, presence of diabetes, smoking history and previous treatment for HTN or dyslipidaemia, or aspirin therapy to estimate the 10-year risk of ASCVD events.³⁷ The presence of more CVD risk-enhancing factors should prompt earlier screening of patients. The AHA defines these risk factors as a family history of premature ASCVD (males <55 and females <65), primary hypercholesterolaemia, metabolic syndrome, chronic kidney disease, inflammatory conditions such as rheumatoid arthritis, lupus or HIV/AIDS, a history of premature menopause (before age 40 years) and a history of APOs that increase lifetime ASCVD risks. Other risk factors such as pre-eclampsia, high-risk race or ethnicity (South Asian ancestry), persistently elevated lipids or biomarkers such as TGs, high-sensitivity C-reactive protein or lipoprotein(a) and/or apolipoprotein B can also increase ASCVD risk and should be accounted for.^{38,39} The USPSTF denotes similar risk factors that should prompt earlier screening, such as a history of diabetes, HTN, smoking, obesity and a family history of CVD.⁴⁰ The current screening recommendations from the AHA and USPSTF are outlined in Figure 2.^{31,32}

In a study of adults in the USA using National Health and Nutrition Examination Survey data, 55% of women aged 20–44 and 74.2% of women older than 45 were considered USPSTF-at-risk with one or more CVD risk factors. Nearly all reproductive age and older women were AHA-at-risk (99.5 and 99.6%). In the same sample, 96.8% of women had seen a healthcare professional within the previous 3 years, but only 62.1% reported having a cholesterol test in the past 5 years.⁵ This suggests that while women maintain their recommended annual health maintenance, they are not appropriately screened for dyslipidaemia. Many clinicians use risk calculators to estimate the 10-year risk of ASCVD, and newer iterations of these calculators, notably the AHA PREVENT calculator, factor in kidney disease and even geographic location as predictors of future CVD events.⁴¹ However, there is not currently a calculator that considers

peripartum complications such as pre-eclampsia or gestational HTN that have been associated with increased lifetime risk; many women's risk is likely undervalued.³⁹

The ACC estimates that eight out of every ten women of childbearing age have never had a lipid assessment.⁴² Earlier screening for dyslipidaemias could be part of perinatal care. This would vastly increase the number of women who are screened. There have also been reports of a correlation between dyslipidaemia, preterm birth and low gestational weight with lifetime risk for CVD. In one study of 1,112 women who were screened at a postpartum visit for dyslipidaemia, 7% had elevated TC, and 13% had elevated TGs. The sample contained 5,101 women, and factors that were associated with screening were race (White), older age and having a diagnosis of coronary artery disease (CAD), HTN and diabetes.⁴³ Identifying dyslipidaemias early in life helps patients and physicians focus on risk modification and early initiation of pharmacotherapy to reduce clinically significant CVD.

Sex disparities in diagnosis and treatment

An estimated 15.5 million Americans over the age of 20 have CVD, and the prevalence increases every year, with a projected 18% increase by 2030.⁴⁴ The 2018 AHA/ACC cholesterol guidelines define dyslipidaemias as high TC (TC >240 mg/dL), high LDL-C (LDL-C >100 mg/dL) and/or high TGs (TG >150 mg/dL).³⁴ Early identification and treatment are the hallmarks of primary and secondary prevention of CVD.^{34,45} The current recommendation is to use a combination of a healthy diet and exercise with high-intensity or maximally tolerated statin therapy to reduce LDL-C by at least 50% in patients with clinical CVD. Statins are the primary recommended pharmacologic treatment in patients who have clinically evident ASCVD, severe hypercholesterolaemia with an LDL-C >190 mg/dL, and diabetes in adults aged 40–75 years.³⁴ A fourth group where statin therapy is recommended is primary prevention in groups that PCEs estimate as being high risk (≥20%), intermediate risk (≥7.5% to <20%) or borderline risk (5% to <7.5%) based on risk enhancers, presence of elevated coronary artery calcium score and

Figure 2: A comparison of current AHA/ACC and USPSTF recommendations for lipid screening^{31,32}

	AHA/ACC	USPSTF
Screening age	20 years Earlier if family history/risk factors	40–75 years Men 35–65 and Women 45–65 with risk factors
Risk factors to prompt early screening	HTN, DM, obesity, family history of premature CVD, smoking, sedentary lifestyle, premature menopause, APOs, CKD, inflammatory conditions, HIV/AIDS, elevated hsCRP or Lp(a), high risk race	Obesity, HTN, smoking, family history of early CVD, metabolic syndrome
Frequency of screening	Every 4–6 years for adults without risk factors Men >45 and Women >55 with risk factors should be screened more frequently Annual screening for confirmed CVD	4–6 years for individuals >40 years without risk factors
Screening factors specific to women	Premature menopause, APOs	None

ACC = American College of Cardiology; AHA = American Heart Association; APOs = adverse pregnancy outcomes; CVD = cardiovascular disease; DM = diabetes; HTN = hypertension; USPSTF = United States Preventive Services Task Force; CKD = chronic kidney disease; hsCRP = high-sensitivity C-reactive protein; Lp(a) = lipoprotein (a)

clinician–patient discussion.¹⁸ The ACC outlined these recommendations in 2019, which are summarized in *Figure 3*.^{18,38} There are no differences in recommendations based on biological sex in the current ACC/AHA guidelines.


In a sample of nearly 60,000 age- and sex-matched Spanish patients included in the ESCARVAL-GENERO study, 18% met the diagnostic criteria for dyslipidaemia but did not have an associated diagnosis or receive prescribed treatment. Female sex was significantly associated with this diagnostic inertia, with the proportion of women fitting this description being higher than that of men (20.1% compared to 15.8%).⁴⁶ In a similar study of the Patient and Provider Assessment of Lipid Management (PALM) registry, 5,693 adults were recommended for statin therapy, with 26.5% not on treatment. Of those patients, 59.2% reported never being offered a statin. Those patients were more likely to be female (odds ratio [OR], 1.22), Black adults (OR, 1.48) and those without insurance (OR, 1.38).⁴⁷ Factors that have been shown in other studies to be positively associated with screening are women who are White with a prior history of CAD, HTN or diabetes.^{48,49} In a sample of 600,000 patients containing 41.7% women in the USA with established risk factors for CVD, 49.9% were not on any statin therapy, and those patients were more likely to be female with higher comorbidities.⁵⁰ Additional studies have been completed comparing statin prescriptions for Black and White women, with Black women having much lower odds of being prescribed statin therapy.⁵¹

Lifestyle modification and aerobic exercise are the initial recommendations listed in many CVD guidelines for dyslipidaemia.⁵² This is supported by the INTERHEART study, which assessed modifiable risk factors such as smoking cessation, vegetable and fruit consumption, and regular physical activity, where intervention reduced the risk of MI by 80%.⁴ While lifestyle modification is essential, it remains unclear whether

this benefit is equal between sexes, and it appears to be more beneficial to men. In the ‘Prevención con Dieta Mediterránea’ or PREDIMED trial (ISRCTN35739639), lifestyle changes, such as adopting a Mediterranean diet, were initially shown to reduce cardiovascular events (myocardial infarction, stroke or death from cardiovascular causes).⁵³ When separating the control versus intervention group by sex, the female subgroup showed no endpoint difference compared with the control, and the publication was later retracted.⁵⁴

If the first tier of recommendations has not been shown to benefit women, it would be expected to see more female patients on statin medications. The opposite is true. In many studies of prescriber behaviour and dyslipidaemia treatment, women are less likely to be prescribed a statin than men, especially younger women.^{50,55,56} In the GOULD observation registry of patients receiving lipid-lowering therapy, women are also less likely to be offered intensification of statin therapy or achieve an LDL <70 mg/dL goal compared to men (23.2% versus 32.4%). This is further compounded by low rates of intensification of lipid-lowering therapy for people of non-White race and Hispanic ethnicity.⁵⁷

Regarding other lipid-lowering medications, women are less likely to have intensification or additional drug additions, even though, as mentioned above, they are less likely to achieve the LDL-C goal. Some drugs have been shown to have preferential benefits for women in clinical trials. For example, the IMPROVED Reduction of Outcomes: Vytorin Efficacy International Trial or IMPROVE-IT trial (NCT00202878), which tested ezetimibe, showed that it reduced LDL-C equally in men and women; however, it significantly lowered the incidence of adverse cardiac events by 12% in women but only by 5% in men.^{58,59} In another recent study of four phase III clinical trials of the drug bempedoic acid, an inhibitor of ATP citrate lyase which works upstream of HMG-CoA enzyme that is affected by statins, women were shown to be 1.6 times more likely than men

Figure 3: ACC recommendations from 2019 for the initiation of statin medication in women^{18,38}

Green box: Statin therapy generally indicated. Red box: statin generally not recommended. Yellow box: statin may be considered.

Risk enhancers include APOs and premature menopause; clinicians should remember to take a comprehensive obstetrical history and discuss any potential risk factors that would indicate the need for statins for primary prevention.

APOs = adverse pregnancy outcomes; ASCVD = atherosclerotic cardiovascular disease; LDL-C = low-density lipoprotein cholesterol; FH = familial hyperlipidaemia

to achieve a minimum LDL-C reduction of 30% in 12 weeks compared to placebo.⁶⁰ While these drugs have been shown to lower LDL-C in women and are effective therapies to achieve LDL-C goals, few women are escalated to these therapies in combination with statins. Since 2018, guidance has been published by the ACC/AHA recommending non-statin medications be used in combination with statin therapy to achieve LDL-C goals.³⁴ There are currently no diagnostic or treatment guidelines that differentiate patients based on biological sex.

Several factors have been hypothesized as reasons for the lack of recognition and treatment of CVD in women. Subpar screening of at-risk women was previously discussed and is a significant factor in the lack of primary prevention initiation. Women are also more likely to have atypical coronary syndromes and have missed or delayed MI diagnoses. This leads to a decrease in the use of statins in secondary prevention and a reduction in overall statin use and intensification.^{52,61} Statin therapy remains the mainstay of treatment for secondary MI prevention, even in women with MI with non-obstructive coronary arteries.^{52,62} Another factor is the teratogenicity of statin therapy and the hesitance to use it in women of childbearing age, although this is no longer a current recommendation of the FDA.²⁶ These factors alone do not explain the lack of statin therapy or LDL-C goal achievement in women. The initiation of statin therapy is driven in large part by PCEs, which often underestimate CVD risk for younger patients, particularly women.^{52,63} Women with high LDL-C levels might have a deferral of

pharmacotherapy as their estimated 10-year risk of MI or stroke is low. However, as previously mentioned, many factors that increase the risk of CVD are not accounted for in PCEs. Additionally, elevated HDL-C levels provide false reassurance in women against the development of future CVD when high LDL-C levels are present; clinicians should instead use a combination of risk factors and PCEs to guide equal and early treatment.^{64,65} A recent recommendation by the ACC/AHA was to use the above risk-enhancing factors with existing calculators to create a more detailed risk profile.⁶⁶

When women are offered statin therapy, they are more likely to decline or distrust the medication, not believing it to be safe or effective.⁵⁵ Studies of statin use in the USA also found that women are more likely to report statin-associated muscle symptoms and thus discontinue therapy.⁶⁷ Another found that women were more likely to try three or more statins but less likely to be treated with alternative lipid-lowering medications. Overall, the study found women to be more dissatisfied with their statin medication and the way their provider explained the treatment, leading to non-compliance.⁸ This suggests that not only do women experience a higher risk of side effects from statins, but there is also a lack of sufficient counselling from providers about the importance of statin therapy. Clinicians must be comfortable with counselling on future CVD risks and use shared decision-making with patients to increase buy-in for treatment.

Future lens: Areas of improvement

In previous sections, we have highlighted several strategies to increase the screening, diagnosis and treatment of women with dyslipidaemia. Still, as more therapies for reducing cholesterol are developed, evaluating treatment groups for these research trials is essential. In several landmark trials, including REDUCE-IT (Evaluation of the Effect of AMR101 on Cardiovascular Health and Mortality in Hypertriglyceridemic Patients With Cardiovascular Disease or at High Risk for Cardiovascular Disease: REDUCE-IT [Reduction of Cardiovascular Events With EPA - Intervention Trial]; ClinicalTrials.gov identifier: NCT01492361), FOURIER (A Double-blind, Randomized, Placebo-controlled, Multicenter Study Assessing the Impact of Additional LDL-Cholesterol Reduction on Major Cardiovascular Events When Evolocumab [AMG 145] is Used in Combination With Statin Therapy In Patients With Clinically Evident Cardiovascular Disease; ClinicalTrials.gov identifier: NCT01764633) and IMPROVE-IT (A Multicenter, Double-Blind, Randomized Study to Establish the Clinical Benefit and Safety of Vytorin [Ezetimibe/Simvastatin Tablet] vs Simvastatin Monotherapy in High-Risk Subjects Presenting With Acute Coronary Syndrome [IMPROVED Reduction of Outcomes: Vytorin Efficacy International Trial - IMPROVE IT]; ClinicalTrials.gov identifier: NCT00202878), which significantly impacted guideline-directed medical therapy for dyslipidaemia, the proportion of women in these trials was less than 30% (29, 25 and 24%, respectively).^{59,68-72} In trials from 1990 to 2018 involving lipid-lowering agents and overall enrolment of women, the average participation rate is only 29%.⁷³ That number drops even further when looking at the inclusion of minority groups; one meta-analysis of the 35 trials of cardiometabolic drugs over the previous 10 years found that only 4% of participants were Black, 12% Asian and 11%


Hispanic, with no increasing trends observed.⁷⁴ This suggests that many lipid-lowering medications are not being tested on a large number of diverse women, and their true potential or side effects are not accounted for. Trials usually exclude women of childbearing age, but women are still underrepresented, given the prevalence of dyslipidaemia. It is paramount to test new lipid-lowering medication adequately in women so that providers can be assured of drug effectiveness.⁶⁴

Additionally, few, if any, of the principal investigators on these clinical trials were women, with a recent analysis showing that only 10.1% of clinical trial committee members were women, and more than half of the trials had no women in clinical trial committee roles.⁷⁵ Another study of over 100 heart failure trials conducted between 2000 and 2019 revealed that only 11% of trial leadership positions were held by women, and this number had not increased significantly over the last 20 years.⁷⁶ Compared to trials led by men, women-led studies were more likely to enrol a higher proportion of female participants (37.9% compared to 44.9%). This suggests that increasing the number of female principal investigators may lead to a greater representation of female study participants and more accurate results from lipid-modifying therapies in female patients. A potential pipeline of increased female participation in clinical cardiology research is outlined in Figure 4.⁷⁶

Conclusion

As discussed, dyslipidaemia and CVD remain a significant concern for women in the 21st century, and the number of female patients with CVD is expected to increase over the next 20 years.^{3,64} The medical community has been historically poor at screening women, starting lipid-lowering

Figure 4: A potential pipeline to increase female representation in clinical trials



Trials with female principal investigators and in committee leadership roles recruit more female participants.⁷⁶ Figure represents a potential pipeline for increasing female participants in research studies

therapy, achieving LDL-C goals and including women in clinical trials. This article aims to draw attention to an issue that remains a key concern in women's health. Increasing awareness of the current state of screening and pharmacological therapy will ideally bring this to clinicians' attention as we work to improve these numbers. Additionally, we hope that

with this increased visibility of the disparities we have outlined above, special attention is brought to the design of clinical trials concerning sex representation and female representation in leadership roles to assess pharmacological interventions for CVD in women adequately. □

1. Garcia M, Mulvagh SL, Merz CNB, et al. Cardiovascular disease in women: Clinical perspectives. *Circ Res*. 2016;118:1273–93. DOI: 10.1161/CIRCRESAHA.116.307547.
2. Mosca L, Benjamin EJ, Berra K, et al. Effectiveness-based guidelines for the prevention of cardiovascular disease in women—2011 update. *Circulation*. 2011;123:1243–62. DOI: 10.1161/CIR.0b013e31820faaf8.
3. Centers for Disease Control and Prevention. NHANES Questionnaires, Datasets, and Related Documentation. Available at: www.cdc.gov/nchs/nhanes/ (accessed: 24 November 2024).
4. Yusuf S, Hawken S, Ounpuu S, et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): Case-control study. *Lancet*. 2004;364:937–52. DOI: 10.1016/S0140-6736(04)17018-9.
5. Robbins CL, Dietz PM, Cox S, et al. Cholesterol screening for women: Who is “at-risk”? *J Womens Health*. 2013;22:404–11. DOI: 10.1089/jwh.2012.4074.
6. Virani SS, Woodard LD, Ramsey DJ, et al. Gender disparities in evidence-based statin therapy in patients with cardiovascular disease. *Am J Cardiol*. 2015;115:21–6. DOI: 10.1016/j.amjcard.2014.09.041.
7. Chou AF, Scholle SH, Weisman CS, et al. Gender disparities in the quality of cardiovascular disease care in private managed care plans. *Womens Health Issues*. 2007;17:120–30. DOI: 10.1016/j.whi.2007.03.002.
8. Karalis DG, Wild RA, Maki KC, et al. Gender differences in side effects and attitudes regarding statin use in the Understanding Statin Use in America and Gaps in Patient Education (USAGE) study. *J Clin Lipidol*. 2016;10:833–41. DOI: 10.1016/j.jacl.2016.02.016.
9. Wild R, Weedon EA, Wilson D. Dyslipidemia in pregnancy. *Endocrinol Metab Clin North Am*. 2016;45:55–63. DOI: 10.1016/j.ecl.2015.09.004.
10. Albrektsen G, Wilsaard T, Heuch I, et al. Lipid levels during adult lifetime in men and women with and without a subsequent incident myocardial infarction: A longitudinal analysis of data from the Tromsø Study 1974 to 2016. *J Am Heart Assoc*. 2023;12:e030010. DOI: 10.1161/JAHA.122.030010.
11. Lührink IK, Wiegman A, Kusters DM, et al. 20-year follow-up of statins in children with familial hypercholesterolemia. *N Engl J Med*. 2019;381:1547–56. DOI: 10.1056/NEJMoa1816454.
12. Expert Panel on Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents and National Heart, Lung, and Blood Institute. Expert panel on integrated guidelines for cardiovascular health and risk reduction in children and adolescents: Summary report. *Pediatrics*. 2011;128(Suppl 5):S213–56. DOI: 10.1542/peds.2009-2107C.
13. Mumford SL, Dasharathy S, Pollack AZ, et al. Variations in lipid levels according to menstrual cycle phase: Clinical implications. *Clin Lipidol*. 2011;6:225–34. DOI: 10.2217/clp.11.9.
14. Jacobson TA, Maki KC, Orringer CE, et al. National Lipid Association recommendations for patient-centered management of dyslipidemia: Part 2. *J Clin Lipidol*. 2015;9:S1–122. DOI: 10.1016/j.jacl.2015.09.002.
15. Yasmin A, Roychoudhury S, Paul Choudhury A, et al. Polycystic ovary syndrome: An updated overview foregrounding impacts of ethnicities and geographic variations. *Life (Basel)*. 2022;12:1974. DOI: 10.3390/12121974.
16. Osibogun O, Ogunmoroti Q, Michos ED. Polycystic ovary syndrome and cardiometabolic risk: Opportunities for cardiovascular disease prevention. *Trends Cardiovasc Med*. 2020;30:399–404. DOI: 10.1016/j.tcm.2019.08.010.
17. Okoth K, Chandan JS, Marshall T, et al. Association between the reproductive health of young women and cardiovascular disease in later life: Umbrella review. *BMJ*. 2020;371:m3502. DOI: 10.1136/bmj.m3502.
18. Cho L, Davis M, Elgendy I, et al. Summary of updated recommendations for primary prevention of cardiovascular disease in women: JACC state-of-the-art review. *J Am Coll Cardiol*. 2020;75:2602–18. DOI: 10.1016/j.jacc.2020.03.060.
19. Magnusson EB, Vatten LJ, Lund-Nilsen TI, et al. Prepregnancy cardiovascular risk factors as predictors of pre-eclampsia: Population based cohort study. *BMJ*. 2007;335:978. DOI: 10.1136/bmj.39366.416817.BE.
20. Mendelson MM, Lyass A, O'Donnell CJ, et al. Association of maternal prepregnancy dyslipidemia with adult offspring dyslipidemia in excess of anthropometric, lifestyle, and genetic factors in the framingham heart study. *JAMA Cardiol*. 2016;1:26–35. DOI: 10.1001/jamacardio.2015.0304.
21. Sharma J, McAlister J, Aggarwal NR, et al. Evaluation and management of blood lipids through a woman's life cycle. *Am J Prev Cardiol*. 2022;10:100333. DOI: 10.1016/j.ajpc.2022.100333.
22. Wild R, Feingold KR. Effect of pregnancy on lipid metabolism and lipoprotein levels. In: Feingold KR, Ahmed SF, Anawalt B, et al. (eds). *Endotext*. South Dartmouth (MA): MDText.com, Inc. 2000.
23. Wiznitzer A, Mayer A, Novack V, et al. Association of lipid levels during gestation with preeclampsia and gestational diabetes mellitus: A population-based study. *Am J Obstet Gynecol*. 2009;201:482. DOI: 10.1016/j.ajog.2009.05.032.
24. Agarwala A, Dixon DL, Gianos E, et al. Dyslipidemia management in women of reproductive potential: An expert clinical consensus from the national lipid association. *J Clin Lipidol*. 2024;18:e664–84. DOI: 10.1016/j.jacl.2024.05.005.
25. Pollack PS, Shields KE, Burnett DM, et al. Pregnancy outcomes after maternal exposure to simvastatin and lovastatin. *Birth Defects Res A Clin Mol Teratol*. 2005;73:888–96. DOI: 10.1002/bdra.20181.
26. US Food and Drug Administration. FDA requests removal of strongest warning against using cholesterol-lowering statins during pregnancy; still advises most pregnant patients should stop taking statins. 2024. Available at: www.fda.gov/drugs/drug-safety-and-availability/fda-requests-removal-strongest-warning-against-using-cholesterol-lowering-statins-during-pregnancy (accessed: 18 January 2025).
27. De Jesus SM, Myerson MM, Gianos EM. Women's Health: Guidelines for Lipid-Lowering in Women – What Have We Learned? Available at: www.lipid.org/lipid-spin/spring-2020/women%E2%80%99s-health-guidelines-lipid-lowering-women%E2%80%93what-have-we-learned (accessed: 15 August 2025).
28. Tan SYT, Teh SP, Kaushik M, et al. Hypertriglyceridemia-induced pancreatitis in pregnancy: Case review on the role of therapeutic plasma exchange. *Endocrinol Diabetes Metab Case Rep*. 2021;2021:21-0017. DOI: 10.1530/EDM-21-0017.
29. Ragunind PF, Cardona I, Muka T, et al. Does reproductive stage impact cardiovascular disease risk factors? Results from a population-based cohort in Lausanne (CoLaus study). *Clin Endocrinol*. 2022;97:568–80. DOI: 10.1111/cen.14730.
30. Ambikairajah A, Walsh E, Cherbuin N. Lipid profile differences during menopause: A review with meta-analysis. *Menopause*. 2019;26:1327–33. DOI: 10.1097/GME.0000000000001403.
31. Henderson KD, Bernstein L, Henderson B, et al. Predictors of the timing of natural menopause in the Multiethnic Cohort Study. *Am J Epidemiol*. 2008;167:1287–94. DOI: 10.1093/aje/kwn046.
32. Honigberg MC, Zekavat SM, Aragam K, et al. Association of premature natural and surgical menopause with incident cardiovascular disease. *JAMA*. 2019;322:2411–21. DOI: 10.1001/jama.2019.19191.
33. Donato GB, Fuchs SC, Oppermann K, et al. Association between menopause status and central adiposity measured at different cutoffs of waist circumference and waist-to-hip ratio. *Menopause*. 2006;13:280–5. DOI: 10.1097/01.gme.0000177907.32634.ae.
34. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APHA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: A report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. 2019;139:e1082–143. DOI: 10.1161/CIR.0000000000000625.
35. Mangione CM, Barry MJ, Nicholson WK, et al. Statin use for the primary prevention of cardiovascular disease in adults: US preventive services task force recommendation statement. *JAMA*. 2022;328:746–53. DOI: 10.1001/jama.2022.13044.
36. United States Preventive Services Task Force. Archived: Lipid Disorders in Adults (Cholesterol, Dyslipidemia): Screening. 2013. Available at: www.uspreventiveservicestaskforce.org/uspstf/recommendation/lipid-disorders-in-adults-cholesterol-dyslipidemia-screening-2008 (accessed: 24 November 2024).
37. Preiss D, Kristensen SL. The new pooled cohort equations risk calculator. *Can J Cardiol*. 2015;31:613–9. DOI: 10.1016/j.cjca.2015.02.001.
38. Arnett DK, Blumenthal RS, Albert MA, et al. 2019 ACC/AHA guideline on the primary prevention of cardiovascular disease: Executive summary: A report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. 2019;140:e563–95. DOI: 10.1161/CIR.0000000000000677.
39. Roeters van Lennep JE, Tokgözoğlu LS, Badimon L, et al. Women, lipids, and atherosclerotic cardiovascular disease: A call to action from the European Atherosclerosis Society. *Eur Heart J*. 2023;44:4157–73. DOI: 10.1093/eurheartj/ehad472.
40. Chou R, Cantor A, Dana T, et al. Statin use for the primary prevention of cardiovascular disease in adults: Updated evidence report and systematic review for the US Preventive Services Task Force. *JAMA*. 2022;328:754–71. DOI: 10.1001/jama.2022.12138.
41. Khan SS, Matsushita K, Sang Y, et al. Development and validation of the American Heart Association's PREVENT equations. *Circulation*. 2024;149:430–49. DOI: 10.1161/CIRCULATIONAHA.123.067626.
42. American College of Cardiology. Women of Childbearing Age Have Staggeringly Low Rates of Lipid Screening. 2019. Available at: www.acc.org/About-ACC/Press-Releases/2019/03/06/10/43/Women-of-Childbearing-Age-Have-Staggeringly-Low-Rates-of-Lipid-Screening (accessed: 20 March 2025).
43. Gopal DJ, Smith CL, Adusumalli S, et al. Screening for hyperlipidemia in pregnant women: An underutilized opportunity for early risk assessment. *J Am Coll Cardiol*. 2019;73:14. DOI: 10.1016/S0735-1097(19)33776-3.
44. Mozaffarian D, Benjamin EJ, Go AS, et al. Heart disease and stroke statistics—2015 update: A report from the American Heart Association. *Circulation*. 2015;131:e29–322. DOI: 10.1161/CIR.000000000000152.
45. Birtcher KK, Ballantyne CM. Measurement of cholesterol. *Circulation*. 2004;110:e296–7. DOI: 10.1161/01.CIR.0000141564.89465.4E.
46. Soriano-Maldonado C, Lopez-Pineda A, Orozco-Beltran D, et al. Gender differences in the diagnosis of dyslipidemia: ESCARVAL-GENERO. *Int J Environ Res Public Health*. 2021;18:12419. DOI: 10.3390/ijerph182312419.
47. Bradley CK, Wang TY, Li S, et al. Patient-reported reasons for declining or discontinuing statin therapy: Insights from the PALM registry. *J Am Heart Assoc*. 2019;8:e011765. DOI: 10.1161/JAHA.118.011765.
48. Kalra DK. Bridging the racial disparity gap in lipid-lowering therapy. *J Am Heart Assoc*. 2021;10:e019533. DOI: 10.1161/JAHA.120.019533.
49. Fan AZ, Greenland KJ, Dai S, et al. Disparities in screening for and awareness of high blood cholesterol—United States, 1999–2002. *JAMA*. 2005;293:1849. DOI: 10.1001/jama.293.15.1849.
50. Nelson AJ, Haynes K, Shambhu S, et al. High-intensity statin use among patients with atherosclerosis in the U.S. *J Am Coll Cardiol*. 2022;79:1802–13. DOI: 10.1016/j.jacc.2022.02.048.
51. Jackson EA, Ruppert K, Derby CA, et al. Is race or ethnicity associated with under-utilization of statins among women in the United States: The Study of Women's Health Across the Nation. *Clin Cardiol*. 2020;43:1388–97. DOI: 10.1002/clc.23448.
52. Varghese T, Lundberg G. Lipids in women: Management in cardiovascular disease prevention and special subgroups. *Curr Cardiovasc Risk Rep*. 2019;13:20. DOI: 10.1007/s12170-019-0615-z.
53. Estruch R, Ros E, Salas-Salvadó J, et al. Primary prevention of cardiovascular disease with a Mediterranean diet. *N Engl J Med*. 2013;368:1279–90. DOI: 10.1056/NEJMoa1200303.
54. Phan BAP, Toth PP. Dyslipidemia in women: Etiology and management. *Int J Womens Health*. 2014;6:185–94. DOI: 10.2147/IJWH.S38133.
55. Nanna MG, Wang TY, Xiang Q, et al. Sex differences in the use of statins in community practice. *Circ Cardiovasc Qual Outcomes*. 2019;12:e005562. DOI: 10.1161/CIRCOUTCOMES.118.005562.
56. Zhang X, Ji L, Ran X, et al. Gender disparities in lipid goal attainment among type 2 diabetes outpatients with coronary heart disease: Results from the CCMR-38 study. *Sci Rep*. 2017;7:12648. DOI: 10.1038/s41598-017-13066-z.
57. Cannon CP, de Lemos JA, Rosenson RS, et al. Use of lipid-lowering therapies over 2 years in GOULD, a registry of patients with atherosclerotic cardiovascular disease in the US. *JAMA Cardiol*. 2021;6:1060–8. DOI: 10.1001/jamacardio.2021.1810.
58. Blazing MA, Giugliano RP, Cannon CP, et al. Evaluating cardiovascular event reduction with ezetimibe as an adjunct to simvastatin in 18,144 patients after acute coronary syndromes: Final baseline characteristics of the IMPROVE-IT study population. *Am Heart J*. 2014;168:205–12. DOI: 10.1016/j.ahj.2014.05.004.
59. Kato ET, Cannon CP, Blazing MA, et al. Efficacy and safety of adding ezetimibe to statin therapy among women and men: Insight from IMPROVE-IT (Improved Reduction of Outcomes: Vytorin Efficacy International Trial). *J Am Heart Assoc*. 2017;6:e006901. DOI: 10.1161/JAHA.117.006901.
60. Goldberg AC, Banach M, Catapano AL, et al. Evaluation of the efficacy and safety of bempedoic acid in women and men: Pooled analyses from phase 3 trials. *Atherosclerosis*. 2023;384:117192. DOI: 10.1016/j.atherosclerosis.2023.117192.
61. Lichtman JH, Lefkowitz EC, Safdar B, et al. Sex differences in the presentation and perception of symptoms among young patients with myocardial infarction. *Circulation*. 2018;137:781–90. DOI: 10.1161/CIRCULATIONAHA.117.031650.
62. Thygesen K, Alpert JS, Jaffe AS, et al. Fourth universal definition of myocardial infarction (2018). *J Am Coll Cardiol*. 2018;72:2231–64. DOI: 10.1016/j.jacc.2018.08.1038.
63. Pencina MJ, D'Agostino RB Sr, Larson MG, et al. Predicting the 30-year risk of cardiovascular disease. *Circulation*. 2009;119:3078–84. DOI: 10.1161/CIRCULATIONAHA.108.166694.
64. Patel N, Mittal N, Wilkinson MJ, et al. Unique features of dyslipidemia in women across a lifetime and a tailored approach to management. *Am J Prev Cardiol*. 2024;18:100666. DOI: 10.1016/j.ajpc.2024.100666.
65. Lázaro P, Murga N, Aguilar D, et al. Therapeutic inertia in the outpatient management of dyslipidemia in patients with ischemic heart disease. The inertia study. *Rev Esp Cardiol*. 2010;63:1428–37. DOI: 10.1016/s1885-5857(10)70277-2.
66. Lloyd-Jones DM, Morris PB, Ballantyne CM, et al. 2022 ACC expert consensus decision pathway on the role of nonstatin therapies for LDL-cholesterol lowering in the management of atherosclerotic cardiovascular disease risk: A report of

- the American College of Cardiology Solution Set Oversight Committee. *J Am Coll Cardiol*. 2022;80:1366–418. DOI: 10.1016/j.jacc.2022.07.006.
67. Navar AM, Peterson ED, Li S, et al. Prevalence and management of symptoms associated with statin therapy in community practice. *Circ Cardiovasc Qual Outcomes*. 2018;11:e004249. DOI: 10.1161/CIRCOUTCOMES.117.004249.
68. Bhatt DL, Steg PG, Brinton EA, et al. Rationale and design of REDUCE-IT: Reduction of cardiovascular events with icosapent ethyl-intervention trial. *Clin Cardiol*. 2017;40:138–48. DOI: 10.1002/clc.22692.
69. Sabatine MS, Giugliano RP, Keech AC, et al. Evolocumab and clinical outcomes in patients with cardiovascular disease. *N Engl J Med*. 2017;376:1713–22. DOI: 10.1056/NEJMoa1615664.
70. Cannon CP, Blazing MA, Giugliano RP, et al. Ezetimibe added to statin therapy after acute coronary syndromes. *N Engl J Med*. 2015;372:2387–97. DOI: 10.1056/NEJMoa1410489.
71. Sever P, Gouni-Berthold I, Keech A, et al. LDL-cholesterol lowering with evolocumab, and outcomes according to age and sex in patients in the FOURIER trial. *Eur J Prev Cardiol*. 2021;28:805–12. DOI: 10.1177/2047487320902750.
72. Bhatt DL, Steg PG, Miller M, et al. Cardiovascular risk reduction with icosapent ethyl for hypertriglyceridemia. *N Engl J Med*. 2019;380:11–22. DOI: 10.1056/NEJMoa1812792.
73. Khan SU, Khan MZ, Raghu Subramanian C, et al. Participation of women and older participants in randomized clinical trials of lipid-lowering therapies: A systematic review. *JAMA Netw Open*. 2020;3:e205202. DOI: 10.1001/jamanetworkopen.2020.5202.
74. Khan MS, Shahid I, Siddiqi TJ, et al. Ten-year trends in enrollment of women and minorities in pivotal trials supporting recent US Food and Drug Administration approval of novel cardiometabolic drugs. *J Am Heart Assoc*. 2020;9:e015594. DOI: 10.1161/JAHA.119.015594.
75. Denby KJ, Szpakowski N, Silver J, et al. Representation of women in cardiovascular clinical trial leadership. *JAMA Intern Med*. 2020;180:1382–3. DOI: 10.1001/jamainternmed.2020.2485.
76. Eliya Y, Whitelaw S, Thabane L, et al. Temporal trends and clinical trial characteristics associated with the inclusion of women in heart failure trial steering committees: A systematic review. *Circ Heart Fail*. 2021;14:e008064. DOI: 10.1161/CIRCHEARTFAILURE.120.008064.